

**A STUDY ON "COMPARATIVE STUDY OF PROPHYLACTIC
RETENTION SUTURING VERSUS PRIMARY CLOSURE IN
LAPAROTOMIES FOR PERFORATION PERITONITIS AT
COIMBATORE MEDICAL COLLEGE HOSPITAL"**

A DISSERTATION SUBMITTED TO

THE TAMILNADU DR.M.G.R MEDICAL UNIVERSITY



In partial fulfilment of the regulations for the award of the degree of

M.S. GENERAL SURGERY – BRANCH I



DEPARTMENT OF GENERAL SURGERY

COIMBATORE MEDICAL COLLEGE AND HOSPITAL

THE TAMILNADU DR.M.G.R MEDICAL UNIVERSITY

CHENNAI

MAY 2019

CERTIFICATE

Certified that this is the bonafide dissertation done by
DR. HARIDOSS.R and submitted in partial fulfilment of the requirement for
the Degree of M.S. General Surgery, Branch I of the Tamilnadu Dr.M.G.R.
Medical University , Chennai.

DATE:

UNIT CHIEF

DATE:

PROFESSOR & HOD

DEPARTMENT OF GENERAL SURGERY

DATE:

DEAN

COIMBATORE MEDICAL COLLEGE

COIMBATORE – 641014.



Coimbatore Medical College

COIMBATORE, TAMILNADU, INDIA - 641 014

(Affiliated to The Tamilnadu Dr. MGR Medical University, Chennai)



ETHICS COMMITTEE



Name of the Candidate: **Dr. Haridoss .R**

Course : **MS (General Surgery) Post Graduate**

Period of Study : **1 year**

College : **Coimbatore Medical College & Hospital.**

Dissertation Topic : **Comparitive study of prophylactic retention suturing versus primary closure in laparotomies for perforation peritonitis**

The Ethics Committee, Coimbatore Medical College has decided to inform that your Dissertation Proposal is accepted and you are permitted to proceed with the above Study.

23.12.16


Member Secretary
Ethics Committee

URKUND

Fwd: [Urkund] 2% similarity - x D42516632 - dissertation131 x Home - URKUND D42483770 - A study on clin x coimbatore medical college x

https://secure.urkund.com/view/41547891-574023-432856#q1bKLvayjibQMdQx8BImOivY6fjqWsTpKxZnpeZlpmcmlcmpSIYGegaGFsYmBgZmpgaWhszG...

Sources	Highlights
Document Submitted Submitted by Receiver Message	<p>dissertation131018.docx (D42516632) 2018-10-14 04:11 (+05:0-30) Haridoss (haridoss.tvmc@gmail.com) haridoss.tvmc.mgmvc@analysis.urkund.com Show full message</p> <p>2% of this approx. 26 pages long document consists of text present in 4 sources.</p>
	Alternative sources
	Dr Ketan Thesis.docx PLAGER.docx thesis - Cert.docx Dissertation FINAL-1.pdf

2 Warnings Reset Export Share

Urkund's archive: Tamil Nadu Dr. M.G.R. Medical University / Dissertation FINAL-1.pdf 70%

PAGE NO 1 INTRODUCTION 1.2 AIM AND OBJECTIVES 4.3 REVIEW OF LITERATURE 6.4 MATERIALS AND METHODS 50.5 OBSERVATION AND ANALYSIS 57.6 DISCUSSION 69.7 CONCLUSION 72.8

#1 Active

70%

PAGE NO 1. INTRODUCTION 2. AIMS AND OBJECTIVES 3. METHODOLOGY 4. REVIEW OF LITERATURE 5. OBSERVATION AND ANALYSIS 6. DISCUSSION 7. CONCLUSION 8.

BIBLIOGRAPHY 9. ANNEXURES

PROFORMA

CONSENT FORM

MASTER CHART

INTRODUCTION

INTRODUCTION Acute wound failure (wound dehiscence or a burst abdomen) refers to postoperative separation of the abdominal musculo-aponeurotic layers.

It is among the most dreaded complications faced by surgeons and is of great concern because of the risk of evisceration.

Acute wound failure occurs in approximately 1% to 3% of patients who undergo an abdominal operation. Dehiscence most often develops 7 to 10 days postoperatively but may occur anytime after surgery from 1 to more than 30 days. A multitude of factors may contribute to wound dehiscence.

ilovepdf_merged (3).pdf ilovepdf_merged (2).pdf Dr Priyadharshini....docx Show all

DECLARATION

I solemnly declare that the dissertation titled “**COMPARATIVE STUDY OF PROPHYLACTIC RETENTION SUTURING VERSUS PRIMARY CLOSURE IN LAPAROTOMIES FOR PERFORATION PERITONITIS AT COIMBATORE MEDICAL COLLEGE HOSPITAL**” was done by me from December 2016 to December 2017 under the guidance and supervision of **PROF. DR. A.NIRMALA, M.S, D.G.O.**

This dissertation is submitted to the Tamilnadu Dr.M.G.R. Medical University towards the partial fulfilment of the requirement for the award of M.S.Degree in General Surgery (Branch I).

PLACE:

SIGNATURE OF CANDIDATE

DATE:

ACKNOWLEDGEMENT

I owe my reverential gratitude and humble thanks to Lord God Almighty for all his mercy, for being with me and showering abundant blessing upon me throughout the course of the study.

I am obliged to record my immense gratitude to **DR. B. ASOKAN Mch**, The Dean, Coimbatore Medical College Hospital for providing all the facilities to conduct the studies.

I express my deep sense of gratitude and heart felt thanks to Professor **DR. V. ELANGO, M.S**, Head of Department of General Surgery for his dynamic guidance, constant help and encouragement throughout the study.

I express my respectful gratitude and indebtedness to my guide Professor **DR. A. NIRMALA M.S.,D.G.O**, for her valuable guidance and support.

I would like to express my sincere thanks to, Professor **Dr. S.SARADHA M.S**, Professor **Dr.Balasubramaniam**, Professor **Lekshminarayani**, Professor **Dr. Ganesh Babu** and **Dr.Srinivasan**. I deeply thank **Dr. P.Murugadasan M.S**, **Dr.Balamurugan M.S**, **Dr.Chandraseker M.S.**, assistant professors of surgery, for all the needful help they have provided for the study.

I acknowledge my gratitude to our Registrar **Dr.Ravi M.S** and all my assistant professors of Department of surgery for their encouragement and support.

I am thankful to The ETHICAL COMMITTEE of Coimbatore Medical College for permitting me to proceed with this dissertation.

Lastly , I am grateful to all the patients whose cooperation made this work possible

ABSTRACT

BACKGROUND

Abdominal wound dehiscence either partial or complete, a common complication of laparotomy and causes a significant mortality and morbidity, with prolonged hospital stay and repeated surgical interventions. So if any system to predict possibility of wound dehiscence, prophylactic retention suturing can be done to prevent adverse events. This study is done to compare such Retention suturing with conventional primary closure in Emergency Laparatomies done for perforation peritonitis.

The objective of this study was to assess the reduced rate of dehiscence in midline laparotomy using prophylactic retention sutures in high- risk patients.

MATERIALS AND METHOD

Our study included 60 patients who underwent Emergency laparotomy for perforation peritonitis under regional or general anaesthesia were randomized into two groups of 30 each. The study group with monitoring of wound healing, pain, hospital stay, wound gaping, wound infection, evisceration in post operative period.

RESULTS

In our study, we have derived that , 14 out of 30 patients (46.7%) who underwent convention primary closure developed wound dehiscence when

compared to 3 out of 30(10.0%) in prophylactic retention suturing group . 17 out of 60 patients underwent re-surgery.

Also there was significant difference in the post operative pain($p=0.001$) and duration of hospital stay($p=0.001$) lesser in retention suturing group.

CONCLUSION

Study concludes that Prophylactic Retention suturing in patients with perforation peritonitis undergoing emergency midline laparotomy decreases the incidence of wound dehiscence, reduces pain and lessens hospital stay in high risk patients, when compared with conventional primary wound closure.

KEYWORDS: prophylactic, Retention suturing, perforation peritonitis, wound gaping

TABLE OF CONTENTS

	CONTENTS	PAGE NO
1.	INTRODUCTION	1
2.	AIMS AND OBJECTIVES	3
3.	MATERIALS & METHODS	4
4.	REVIEW OF LITERATURE	7
5.	OBSERVATION AND ANALYSIS	57
6.	DISCUSSION	94
7.	CONCLUSION	97
8.	BIBILOGRAPHY	
9.	ANNEXURES	
	PROFORMA	
	CONSENT FORM	
	MASTER CHART	

INTRODUCTION

Acute wound failure (wound dehiscence or a burst abdomen) refers to postoperative separation of the abdominal musculo-aponeurotic layers. Acute wound failure occurs in approximately 1% to 3% of patients who undergo an abdominal operation. Dehiscence most often develops 7 to 10 days postoperatively but may occur anytime after surgery, from 1 to more than 20 days. A multitude of factors may contribute to wound dehiscence. Acute wound failure is often related to technical errors in placing sutures too close to the edge, too far apart, or under too much tension. Local wound complications such as hematoma and infection can also predispose to localized dehiscence. Increased intra-abdominal pressure (IAP) is often blamed for wound. In healthy patients, the rate of wound failure is similar whether closure is accomplished with a continuous or interrupted technique. In high-risk patients, however, continuous closure is worrisome because suture breakage in one place weakens the entire closure. Predisposing factors which may include Technical error in fascial closure, Emergency surgery, Intra-abdominal infection, Advanced age, Wound infection, hematoma, and seroma, Elevated intra-abdominal pressure, Obesity, Chronic corticosteroid use, Previous wound dehiscence, Malnutrition, Radiation therapy and chemotherapy, Systemic disease (uremia, diabetes mellitus).

Generalized peritonitis patients requiring emergency laparotomies are at high risk of post operative complications including wound infections, dehiscence and evisceration.

In addition poor nutritional status and delayed presentation is associated with increased incidence of wound dehiscence.

Additionally, total cost of treatment increases both in terms of lengthy hospital stay, nursing and man power utility in treating this patient and its complications.

Different surgical techniques /sutures for closing the laparotomy wounds are being advocated such as interrupted or continuous suturing, mass closure/ layered closure , delayed absorbable/non-absorbable . Abdominal wound dehiscence is surgically managed by retention sutures, mesh, biological implant placement, and interrupted X sutures.

There is decreased incidence of wound dehiscence in-patients undergoing laparotomies for perforation peritonitis, by prophylactic placement of retention sutures. Thereby reducing post operative morbidity and mortality and better outcome.

Main objective of the study is to compare the wound healing in prophylactic retention suturing verses conventional primary closure in emergency laparotomies for perforation peritonitis.

Secondarily to compare, incidence of wound dehiscence, wound infection, post operative pain, re-surgery and hospital stay.

AIM AND OBJECTIVE

The aim of the study is to compare the efficacy of prophylactic retention suturing technique versus conventional primary closure in patient undergoing midline laparotomy for perforation peritonitis in terms of wound dehiscence, post operative pain, hospital stay and re-surgery.

MATERIALS AND METHODS

This comparative study of wound healing in perforation peritonitis is based on the patients admitted with signs and symptoms of peritonitis due to gastrointestinal perforation for a period of 12 months from December 2016 to December 2017, in general surgery department of Coimbatore medical college hospital, Coimbatore.

A total of 60 patients presenting with perforation peritonitis at emergency department were subjected to emergency midline laparotomy. They are divided into two groups by simple random sampling.

For all the cases detailed history, complaints, clinical features were taken and vital signs recorded.

These patients were subjected to investigations – complete blood counts, renal function tests, serum electrolytes, liver function tests with serum proteins, electrocardiogram, X-ray chest and abdomen erect, viral markers and blood grouping & Rh typing.

STUDY DESIGN:

Prospective comparative study.

DURATION OF STUDY:

12 months (December 2016 to December 2017)

SAMPLE SIZE:

Simple random sampling

Group A- conventional primary closure done (30).

Group B- prophylactic retention suturing (30).

INCLUSION CRITERIA:

- 1) Patients with features of perforation peritonitis undergoing emergency laparotomy.
- 2) Patients age group 20 years and above.
- 3) Patients with Anaemia.
- 4) Patients with Hyperbilirubinemia.
- 5) Patients with Hypoproteinemia.

EXCLUSION CRITERIA:

- 1) Age less than 20 years.
- 2) Immuno-compromised patient.

METHODOLOGY

After proper clinical assessment the patients were actively resuscitated with analgesics, intravenous fluids, nasogastric aspiration and antibiotics. The bladder was catheterized to monitor the urine output.

After stabilizing the general condition, the patients were taken up for surgery.

Postoperatively nasogastric aspiration was continued, nutrition and electrolyte balance were maintained with intravenous fluids. Daily the patients were assessed for recovery and complaints if any were recorded.

A separate proforma for each case containing all the relevant particulars were maintained. The patients were followed up in the post operative period and the post operative outcomes of all patients were documented and graphed into groups separately.

The incidence of all the post operative outcomes for pain, wound infection, seroma formation, wound dehiscence and evisceration are statically computed as percentage and the final results was given in accordance to the study and observations made. Patients were monitored in the post operative period for pain, wound infection, seroma formation, wound dehiscence, and evisceration. All data were recorded and statistically analysed.

Specific instruction was given to each patient on discharge, to come for periodical review regularly.

REVIEW OF LITERATURE

Intra abdominal infections have been well recognized throughout the history of medicine. Widespread application of sophisticated investigations, prompt surgical intervention and effective antibiotic therapy have significantly reduced the mortality from 90% at the initial part of the century to 10-20 % today. Despite these advances mortality persists, with the patients succumbing to the effect of sepsis and eventual multisystem organ failure.

Peritonitis can be defined as a state where there is inflammation of the peritoneum in response to an injury or the infection. A perforation is one that extends through the wall of the gastrointestinal tract, establishes communication between the lumen of the viscus and the surrounding peritoneal cavity, causing peritonitis.

The initial peritonitis is followed by most dramatic and devastating sequence of events which if not properly managed may cause severe morbidity and mortality. So early detection and prompt definitive treatment which is usually surgical, measures a lot in the prognosis of the patient.

HISTORICAL ASPECTS:

Peritonitis due to perforation of peptic ulcer was described by Littre in 1970- In his case study of acute perforated peptic ulcer, described a lady patient HENNEITTA ANNA – Duchess of Orleans and daughter of CHARLES I. She

developed signs of peritonitis and died within 9 hours of the onset of symptoms. The autopsy revealed a perforated gastric ulcer.

CHRISTOPHER RAWLINSON reported one case in 1729 and JACOB PENADA another in 1795. HANBURGEA in 1746 presented the first duodenal ulcer perforation. In a review of literature, the clinical features of peritonitis was described by CRUEILEIR in 1835. Illustrations of peptic ulcer perforation were recorded in his Atlas. BRINTON made a collection of 234 cases in 1857. HEUSNER was first surgeon to perform a simple closure of perforated ulcer in 1892. Duodenal ulcer perforation closure was done first by DEANAM in 1894.

In 1929 CELLON JONES developed the technique of using live omental patch. Contributions in this field came from ROSCOE GRAHAM- 1937 developed technique of free omental patch. Perforative peritonitis was a rare disease some 100 years ago. RODNEY MAINGOT reports that 30 years ago the mortality in perforation was 30%. Ten years later it has declined to 10% ILLINGWORTH and AVERY JONES et.al found that the mortality was 5 % in 1953 and now it is less than 5%.

Contributions in this field came from ROSCOE GRAHAM- 1937

LANGMAN- 1979

WALT & COLLOGNES- 1982

BONNERIE- 1985

ANATOMY

ANATOMY OF PERITONEUM

James Douglas of Edinburgh first described the peritoneal cavity in 1730. Peritoneum is a serous membrane lining the peritoneal cavity. It invests the abdominal structures. Peritoneum consists of visceral and parietal layers. It is divided in greater sac and lesser sac communicating through foramen of Winslow. Peritoneum has got the surface area of two meters normally it contains about 75-100 ml of clear straw-coloured fluid.

PERITONEAL FOSSAE

The right and left sub-phrenic spaces lie between diaphragm and liver and are on each side of the Falciform Ligament. The hepato-renal recess or pouch lies between the right lobe of liver, Right kidney and right colic flexure. These lie above the greater omentum.

Duodenal fossae, caecal fossae and pelvic fossae are below the greater omentum. In the male the Recto-vesical fossae lies between rectum and bladder. In the female the uterus and its broad ligaments divide this recto-vesical pouch into vesico uterine and Recto- uterine fossae.

THERE ARE FOUR GUTTERS

1. The right lateral gutter is placed lateral to the ascending colon and caecum.

2. The left lateral gutter is placed lateral to the descending colon and sigmoid.
3. The right medial gutter lies between the root of the mesentry and ascending colon.
4. The left medial gutter lies between the root of the mesentry and descending colon.

Right medial gutter is closed above and below. Other three lead to the Pelvis. The right lateral gutter conducts from the right sub-phrenic space and hepatorenal pouch into the pelvis in sitting posture.

SURGICAL PHYSIOLOGY

Peritoneal cavity is the largest cavity in the body, the surface area of its lining membrane is 1.8m^2 , equal to the surface area of the skin. This serous membrane is composed of single layer of polyhedral mesothelial cells resting upon a thin layer of fibroblastic tissue. Beneath these two layers lies a highly vascularised connective tissue with a network of lymphatic tissue.

It is estimated that edema of the peritoneum of 1 mm thickness may lead to sequestration of 18 Ltrs of fluid. It functions as a passive semi permeable membrane to the diffusion of water, electrolytes and macrolutes.

Normally < 50 ml of sterile, pale yellow coloured peritoneal fluid is present in the cavity. It resembles lymph fluid, has low specific gravity (1.000)

and contains lymphocytes and polymorphs. The cell count is usually <3,000 cells per cubic mm.

Peritoneal fluid is secreted by visceral peritoneum, circulated through the peritoneal cavity, finally the fluid is mostly absorbed into the lymphatic circulation via peritoneal surfaces and also through diaphragmatic lymphatic. Negative intra thoracic pressure during inspiration facilitates this fluid movement into thoracic lymph channel.

Bacterial clearance from peritoneal cavity depends on,

- 1) Subdiaphragmatic lymphatic channels
- 2) Phagocytosis by peritoneal macrophages.

These two local mechanisms represent the first line of defense against the bacterial contaminations.

PATHO PHYSIOLOGY

Following any type of injury the peritoneum reacts by way of inflammation the intensity of which varies according to the etiology.

The sequence of events being, the release of Histamine and other permeability promoting factors from the Mast cells, resulting in the permeation and exudation of protein rich plasma, containing fibrinogen into the peritoneal cavity. In addition, Thromboplastin and tissue plasminogen activators are also released which are very much needed for the conversion of fibrinogen into fibrin which is responsible for the adhesion formation.

The healing of the peritoneal defect is extremely fast and in many a times healing is complete by 3 days and mesothelial regeneration is complete by the 8th day.

Unlike the skin which heals from the centripetal growth from the wound margin peritoneal defect is repaired from everywhere simultaneously.

Apart from the peritoneal response other manifestation include the metabolic response, Bowel response and hypovolemia. A surgeon dealing with a case should bear this in mind.

Metabolic response is in short by way of increased glucose conception, increased anaerobic metabolism because of glycolysis.

Bowel reacts to peritonitis initially by increased motility only to gradually become abscent and adynamic ileus results. It is favourable site for accumulation of gas and fluid in the third compartment.

The fluid which accumulates in the gut is mostly from the extra cellular compartment, constituting like that of plasma which ultimately results in hypovolemia.

BACTERIOLOGY

Despite the massive contamination and complexity the microbial spectrum that results from perforative peritonitis within 48 hours, is narrow and only few organisms could be isolated. Among them, commonest are, E.Coli,

Anaerobic Bacteroides species, Enterococci, Aerobic and anaerobic Streptococci and clostridia species.

E.coli and Enterococcus are predominating during peritonitis phase while Bacteroids predominate phases.

Factors favour the localization of peritonitis


- 1) Anatomically the peritoneal cavity proper is subdivided into supracolic and infracolic compartments by the transverse colon and its mesocolon. This decreases the spread of infection from one to the other.
- 2) Formation of fibrinous adhesion between the loops and parietis.
- 3) Outpouring of serous fluid rich in leucocytes and antibodies.
- 4) Paralytic ileus.
- 5) Greater omentum envelops the inflamed structures.

Factors cause the spread of Peritonitis

- 1) When perforation occurs suddenly before protective mechanisms have been mobilized.
- 2) Ingestion of food stimulates the peristaltic activity which favours diffusion.
- 3) Purgatives and enema
- 4) Virulent organisms

- 5) In children – Omentum is small
- 6) Injudicious and rough handling of localized collection.
- 7) Immunodeficiency states- AIDS, Immunosuppressant drugs etc.

**X – RAY CHEST – PA – VIEW-
GAS UNDER THE DIAPHRAGM**



ETIOLOGY

1) TRAUMA

- Blunt
- Penetrating – Incised clean wounds as knife wounds/ Ragged or blunt wounds.
- Iatrogenic

2) PEPTIC ULCER DISEASE

3) INFLAMMATORY

- Diverticulars
- Ulcerative colitis
- Crohn's disease

4) VASCULAR

- Ischaemic colitis
- Volvulus
- Vascular occlusion

5) Mechanical

- Volvulus
- Obstruction
- Band
- Malignant – proximal perforation

6) MALIGNANT PERFORATION - Through site of the lesion

7) BARIUM ENEMA/MEAL

8) FOREIGN BODY

9) SPONTANEOUS PERFORATION

Perforations secondary to peptic ulcer may be either gastric ulcer perforation or duodenal ulcer perforation, which again can be of acute onset or chronic ulcer going in for perforation. The causative reasons for peptic ulcer are beyond the scope of this study but the precipitating causes for perforations are discussed later.

In contrast to the peptic ulcer, trauma is showing itself to be an ascending cause of perforative peritonitis. It can be in the form of penetrating injuries which constitute stab injuries, penetrating Road traffic injuries, Industrial Injuries, Gun shot injuries and less commonly bull gore injuries. Blunt injuries like fall from a height or road traffic accidents produce perforative peritonitis by way near of disruption in continuity of hollow viscus more commonly near the fixed points of the small intestine.

Through inflammatory bowel disease, Typhoid, Tuberculosis and Amoebiasis are very much common, these being the cause of peritonitis is very much less for the prevalence of the disease.

Maligancy is a disease of old age and in particular, gastric malignancy more so ulcerative type occasionally results in perforation.

Ischaemia of the bowel can cause peritonitis either by ischaemic necrosis of the bowel wall resulting in peritonitis, or the free permeation of toxic material and bacteria across the non-viable intact bowel. Both

varieties are seen in cases like Volvulus, strangulated hernias and obstruction due to strictures.

CLINICAL FEATURES

Perforation peritonitis is the most common type of peritonitis. As already discussed the peritoneal infection is arising from an intra abdominal source, usually perforation of a hollow viscus. The majority of these episodes are the results of primary lesions of the Stomach, Duodenum, Small intestine, Colon and Appendix. Depends upon the severity of the lesions peritonitis may be a localized or diffused.

FEATURES OF LOCALISED PERITONITIS

Patients will have abdominal pain and vomiting. Systemic signs like fever and tachycardia will be present. Important signs like guarding and rigidity of the abdominal wall over the inflamed area will be obvious. Each area will have peculiar presentation, for example, shoulder tip pain in subphrenic abscess, urinary symptoms and mucus diarrhoea in pelvic peritonitis. Here in pelvic peritonitis, abdominal signs are less but tenderness on PV or RR is more pronounced.

The localized peritonitis may resolve with appropriate treatment or followed by abscess formation or become diffuse peritonitis.

DIFFUSE (GENERALISED) PERITONITIS

The clinical course can be divided into three phases which overlap with each other:

- 1) Initial phase
- 2) Intermediate phase
- 3) Terminal phase

1. INITIAL PHASE

Abdominal pain is almost always the predominant presenting symptom. It is first experienced at the site of the original lesion and spreads outwards from here. To start with it is a sudden, sharp and severe pain. In fully established peritonitis it is constant, burning and aggravated by movement or breathing. So that the patient remains still and in recumbent position.

Anorexia, Nausea and Vomiting, as well as thirst and oliguria are frequently present. Systemic signs like Fever, Diaphoreses, Tachycardia and Hypotension are usually present.

Abdominal tenderness, guarding and rigidity (referred as board like) are extensive in anterior abdominal wall involvement, and are less in pelvic peritonitis and in lesser sac collection. Obliteration of liver dullness is an important diagnostic sign, but its absence does not rule out the perforation. Bowel sounds are diminished and with the onset of paralytic ileus it become absent.

These typical features may be absent in very young and old aged, immunosuppressed, quadriplegic, comatose patients and in early post operative period.



2. INTERMEDIATE PHASE:

If unattended, the patient goes into this phase. Some times this phase is called as Stage of Delusion, because here the patient feels better with slow pulse, decrease pain and tenderness leaving a silent and soft abdomen which misleads the observer. Some times with this phase the peritonitis may become localized with the formation of abscess.

3. TERMINAL PHASE

If resolution or localization have not occurred the patient develops toxemia and shock which is evident by effortless regurgitation of feculent material, paralytic ileus with increasing distended and silent abdomen. Signs of circulatory failure like cold and clammy extremities, sunken eyes, dry tongue, thread irregular pulse, drawn and anxious face (Hippocratic facies) are obvious. Finally the patients lapses into unconsciousness due to multi system failure.

HOLLOW VISCUS PERFORATION



MANAGEMENT

The primary objectives in the management of secondary peritonitis are,

- 1) Resuscitation and general care of the patient
- 2) Antibiotic therapy
- 3) Surgical management
- 4) Continued metabolic support

1) Resuscitation and general care of the patient:

a) Intravenous fluids

Plasma volume must be restored and the electrolyte concentration also to be corrected along with plasma protein replacement. The effectiveness of the therapy is judged by the normalization of pulse, blood pressure and mental status. Invasive peripheral arterial and central cardiac pressure monitoring catheters (Swan – Ganz catheters) should be placed in patients with specific shock, old age or cardiac, pulmonary, renal insufficiencies.

b) Nasogastric aspiration

Ryles tube is passed into the stomach to decompress it, this prevents pulmonary aspiration and abdominal distension. Intermittent aspiration is maintained till the ileus is resolved.

c) Analgesia

Pain must be relieved before and after the surgery. Morphine may be given and continued for 8 hours. This allows early mobilization and physiotherapy in the postoperative period which prevents complications like atelectasis, deep vein thrombosis and pulmonary embolism.

d) Oxygen is needed especially in septic shock

e) Fluid balance chart

Urinary drainage catheter is essential to monitor the hourly urine output which is a reliable indicator of fluid resuscitation. To the urine output, gastric aspiration and other losses are added then fluid requirement is estimated.

This amount to be replaced in phased manner.

f) BLOOD TRANSFUSION IF NECESSARY

g) Others- H₂ – Blockers, Antacid therapy etc.

2. ANTIBIOTIC THERAPY

Administration of antibiotics prevents multiplication of bacteria and the release of endotoxins. As the infection is usually a mixed one, presumptive therapy which covers the aerobic gram negative rods and anaerobic organisms is started. The agents used against aerobic organisms are Aminoglycosides, II & III generation Cephalosporins, Monobactam, Ampicillin with Sulbactam / Clavulanic acid etc. For anaerobic organisms Metronidazole is used. To

prevent the recurrence of sepsis, therapy should be continued for adequate duration, ie. Till the temperature and cell count become normal.

3. SURGICAL MANAGEMENT

In perforation peritonitis surgical control of the infecting organism is the mainstay of the treatment. Here, only basic essential general surgical concepts are discussed. Specific operations for each organ is dealt with in the later part.

The operative management is primarily directed towards the,

- 1) Control of the source of contamination
- 2) Reduction of the bacterial inoculums.

1. CONTROL OF THE SOURCE OF CONTAMINATION:

Once the patient is fit for anaesthesia and surgery, exploration of the abdomen is carried out usually through a midline incision which gives a wide exposure and access to majority of the peritoneal cavity. The contaminating source is identified and dealt with simple closure of the perforation or resection of the perforated viscus or exclusion of the affected organ.

2. REDUCTION OF THE BACTERIAL INOCULUM

The following procedures are generally avocated,

- a. After the cause has been dealt with, whole peritoneal cavity is explored, the collected fluid is sucked out, debridement and removal of fibrin, blood clot

and necrotic material is done. Copious irrigation of peritoneal cavity with 2 to 3 liters of normal saline.

b. Continuous post operative lavage with the placement of multiple catheters and closed suction drainage. Lavage with crystalloid

c. Planned repeated laparotomy or repeated reexploration with the use of zipper debridement of necrotic materials, drain abscesses and post operative recurrent sepsis have received renewed interest.

PROGNOSIS

With modern treatment perforative peritonitis carries a mortality range from 10% to 40% Mortality for duodenal ulcer and appendicular perforation is usually about 0 to 10%, for intestinal perforation 20 to 40% and for postoperative perforation about 0%. The factors influencing the mortality are,

1. Age of the patient – Greater in the older age group.
2. Time interval between the occurrence of perforation and initiation of treatment – there is approximately a five fold increase in the mortality among the patients who received the treatment after 24 hours compared to the patients who reached within 6 hours.
3. Site of perforation – Mortality is more in colonic perforation.
4. Extent of the diseases.
5. Electrolyte imbalance.

6. Undrained collections.
7. Multisystem breakdown – Renal, cardiac, hepatic and pulmonary insufficiencies.
8. Malignancy, diabetes, etc.,

COMPLICATIONS

All the complications of a severe bacterial infection are possible but the specific complications of peritonitis are as follows:

1) Residual Abscess – Can occupy any of the following sites,

- a) Sub phrenic space
- b) Paracolic gutters
- c) Right iliac fosa
- d) Pelvic cavity

In majority of the cases the abscess resolves with proper antibiotic therapy. If the abscess fails to resolve, it can be managed by one of the following methods:

- a) Ultra sound of C.T. guided aspiration
- b) Percutaneous placement of drainage tube under fluoroscopic or Ultra sound control.
- c) Open drainage
- d) Pelvic abscess is drained through pervaginal or perrectal route.


- 2) Paralytic ileus
- 3) Acute Intestinal Obstruction due to peritoneal adhesion.
- 4) Wound infection – Commonest organisms involved are Staphylococcus aureus, Enterococcus, E. Coli, Pseudomonas, Klebsiella, and Proteus infection is more in intestinal perforation.
- 5) Wound dehiscence and burst abdomen
- 6) Fistula and sinus formation
- 7) Deep vein thrombosis
- 8) Pulmonary infection and atelectasis.

BIOLOGY OF WOUND HEALING


Wound healing is the mechanism whereby, body attempts to restore the integrity of the injured part. This falls far short of tissue regeneration by pluripotent cells .

It is the body's response to tissue injuries, this is an essential and most primitive process common to all multicellular organisms. Here a principal type of cell assumes embryonic features and undergoes migration, divides and then differentiates to produce an extracellular matrix in a seemingly less than optimal environment.

Normal wound healing consists of three overlapping phases



Stages of Wound Healing



Factors influencing healing of a wound

Site of the wound

Structures involved

Mechanism of wounding

Incision

Crush

Crush avulsion

Contamination (foreign bodies/bacteria)

Loss of tissue

Other local factors

Vascular insufficiency (arterial or venous)

Previous radiation

Pressure

Systemic factors

Malnutrition or vitamin and mineral deficiencies

Disease (e.g. diabetes mellitus)

Medications (e.g. steroids)

Immune deficiencies (e.g. chemotherapy, acquired immunodeficiency syndrome (AIDS))

Smoking

Classification of wound closure and healing

Primary intention

Wound edges opposed

Normal healing

Minimal scar

_ Secondary intention

Wound left open

Heals by granulation, contraction and epithelialisation

Increased inflammation and proliferation

Poor scar

_ Tertiary intention (also called delayed primary intention)

Wound initially left open

Edges later opposed when healing conditions favourable

wound surface. Fibroblasts require vitamin C to produce collagen.

The wound tissues formed in the early part of this phase is called granulation tissue. In the latter part, there is an increase in the tensile strength of the wound due to increased collagen, which at first deposited in random fashion. It consists of type III collagen.

The remodelling phase is characterised by maturation of collagen (type I replacing type III until a ratio of 4:1 is achieved). There is a realignment of collagen fibres along the lines of tension, decreased wound vascularity and wound contraction due to fibroblast and myofibroblast activity.

TYPES OF HEALING

1. Primary Intention

Usually surgical incisions heal by primary intention, where the wound edges are brought together (apposed) and when held in a place by mechanical means shortly after injury (adhesive strips, staples & sutures), allowing the wound time to heal and develop enough strength to withstand stress without support.

It is also the way most surgical wounds heal. Typically such wound are created in aseptic conditions with minimal bacterial contamination and a minor amount of tissue damage. They have accurately apposed and sutured wound edges. Epithelialization and contraction have little role in this type of healing.

2. Secondary Intention

When the wound is left opened, healing by secondary intention happens, because of the presence of infection, excessive trauma or skin loss, and the wound edges come together naturally by means of granulation and contractions.

There are 3 main reasons why wound undergo this form of healing :

Wound infection, substantial tissue damage or lack of skin edge apposition. This form of repair is also encountered following ulceration, abscess formation, major superficial wounds or tissue infarction. Healing by secondary intention allows the natural process to occur without surgical closure. Wound contraction may be the most important factor that aids secondary healing.

3. Tertiary Intention or Delayed Primary Closure

Often performed in contaminated wounds, does not retard wound strength. Thus delayed closure may decrease wound morbidity without impairing wound strength.

PHASES OF WOUND HEALING

There are essentially 3 phases of wound healing:

LAG PHASE / INFLAMMATORY OR EXUDATIVE PHASE

- Inflammation of the wound and mobilization of the cells which synthesize granulation tissue.
- Lag phase is entitled not because it is a phase of inactivity in wound repair, but simply because there is no significant increase in the mechanical strength of the wound.

Proliferative or granulation phase

- Granulation tissue was formed in the wound; collagen and mucopolysaccharides are synthesized by the granulation tissue, and there is increase in the mechanical strength of the wound.

Wound contraction (matrix formation) or remodelling phase

- Cells in the wound diminish in number but there is extensive remodelling of wound collagen and further increase in mechanical strength of the wound.

INFLAMMATION / EXUDATIVE PHASE (2-5 DAYS)

It begins immediately upon injury and lasts for few days. Tissue injury may cause disruption of blood vessels and extravasation of blood components. The blood clot re-establishes haemostasis and provides extracellular matrix for cell migration. Platelets not only facilitate the formation of a haemostatic plug but also secrete several mediators of wound healing such as platelet derived growth factor that attract and activate macrophages and fibroblasts. Several vasoactive mediators and chemotactic factors are generated by the coagulation and activated complement pathways, which help in recruiting inflammatory leukocytes to the site of injury.

With in the first 5 – 6 hours after injury, neutrophils enter the wound. Infiltrating neutrophils cleanse the wounded area of foreign particles and bacteria are then extruded with the eschar or phagocytosed by macrophages.

Monocytes then infiltrate the wound by 24 – 48 hours in response to specific chemo attractants such as TGF- β . It becomes activated macrophages, which in turn releases growth factors such as platelet – derived growth factors and vascular endothelial growth factors which initiate the formation of granulation tissue. Macrophages bind to specific proteins of the extracellular matrix by their integrin receptors, an action that stimulates phagocytosis of micro organisms and fragments of extra cellular matrix.

Many important cytokines are released by the macrophages like CSF-1, TNF α , TGF- α , IL-1 and other growth factors, which are necessary for the initiation and propagation of the new tissue formed in the wounds. Thus

macrophages appear to have a pivotal role in the transition between inflammation and repair.

EPITHELIALIZATION

In this period, there is also proliferation of epithelial cells at the epidermal-dermal junction, which migrates towards the midline reforming a thin epidermal layer under the surface of the clot in sutured surgical wounds epithelial migration begins within the first 24 hours of the injury and may be completed as early as 72 hours in healthy individuals

Closure of the wound is not the only function of epithelial cells in the inflammatory phase. The development of techniques in molecular biology has led to unequivocal identification of many uninvolved cytokines. Keratinocytes have been shown to produce GM-CSF, TGF- α and fibroblast proliferation and enhances the production of type I and II collagen mRNA and an angiogenic factor. Thus they help to prepare and promote the next phase of wound healing.

PROLIFERATION OF GRANULATION TISSUE (2 days – 3 weeks)

Several chemotactic, growth and activating factors produced in the inflammatory phase are used in the initiation and development of granulations tissues, which lasts for about 4 – 21 days after injury.

Granulation tissue comprise a loose matrix of fibrin ,fibronectin, collagen and glycosaminoglycans, particularly hyaluronic acid, containing macrophages, fibroblasts and in growing blood vessels. In incisional wounds during proliferation phase the wound begins to gain tensile strength, but at the

sametime it is during this phase that wound dehiscence and evisceration most commonly occurs.

FORMATION OF GRANULATION TISSUE

New stroma, called granulation tissue begins to invade the wound space approximately four days after injury. Numerous new capillaries endow the new stroma with its granular appearance. Macrophages, fibroblasts and blood vessels move into the wound space at the same time. The macrophages provide a continuing source of growth factors necessary to stimulate fibroplasias and angiogenesis; the fibroblasts produce the new extracellular matrix necessary that support cell in growth and blood vessels carry oxygen and nutrients to sustain cells metabolism.

Growth factors especially platelet derived growth factors PDGF and TGF β stimulate the fibroblasts to proliferate and migrate into the wound space. The structural molecules o the newly formed extracellular matrix, termed the provisional matrix, contribute to the formation of granulation tissue by providing a conduit for cell migration.

These molecules include: Fibrin, Fibronectin, Hyaluronic acid

The appearances of fibronectin and the appropriate receptors that bind fibronectin, fibrin or both on fibroblast appear to be the rate limiting step in the formation granulation tissue. The fibroblasts are responsible for the synthesis, deposition and remodelling of the extracellular matrix. Conversely the extra cellular matrix can have a feed back effects on the ability of fibroblasts to remodel.

Cell movement into a blood clot of cross – linked fibrin or into tightly woven extracellular matrix requires an active proteolytic system that can cleave a path for cell migration. A variety of fibroblast derived enzymes including plasminogen activator and collagenases are potential candidates for this task.

After migrating into wound, fibroblasts commence the synthesis of extracellular matrix. The provisional matrix is gradually replaced with a collagenous matrix. Once an abundant collagen matrix has been deposited, the fibroblasts stop producing collagen and the fibroblast rich granulation tissue starts getting replaced by a relatively a cellular scar.

Dysregulation of these processes occurs in the fibrotic disorders such as keloid formation.

NEOVASCULARIZATION

Formation of new blood vessels is necessary to sustain the newly formed granulation tissue. Angiogenesis is a complex process that relies on extracellular matrix on in the wound bed and also migration and mitogenic stimulation of endothelial cells.

Induction of angiogenesis has been attributed to molecules like TGF β , angiogenin, angiotropin and vascular endothelial growth factor. Low oxygen tension and elevated lactic acid may also stimulate angiogenesis. Many of these molecules mentioned above appear to induce angiogenesis by stimulating the production of

- a. Basic fibroblast growth factor – active during first days of repair

- b. Vascular – endothelial cell growth factor – critical during formation of granulation tissue on days 4 through to 7.

MECHANISM

Injury causes destruction of tissue and hypoxia. Angiogenesis factors such as fibroblast growth factors are immediately released from macrophages. Proteolytic enzymes released into the connective tissue degrade extra cellular matrix proteins. Fragments of these proteins recruit peripheral blood monocytes to the site of injury, where they become activated macrophages and release angiogenesis factors. These factors stimulate endothelial cells to release plasminogen activator and procollagenases, which in concert get activated and digest basement membranes.

The fragmentation of the basement membranes allows endothelial cells stimulated by angiogenesis factors to migrate and form new blood vessels at the injured site. Once the wound is filled with granulation tissue, angiogenesis occurs and many of the new blood vessels disintegrate as a result of apoptosis.

MECHANISM OF COLLAGEN FORMATION AND NEOVASCULISATION

WOUND CONTRACTION OR REMODELLING PHASE

During second week of healing, fibroblasts assume a myofibroblasts phenotype characterized by a large bundles of actin containing microfilaments disposed along the plasma membrane of cells and by cell – and cell – matrix linkage.

Granulation tissue begins to be remodelled and its vascularity decreases as the amount of collagen increases. Maturation of the scar occurs over the next few months and is characterized by further remodelling. Collagen produced from fibroblasts is initially laid down in a vertical manner; but gradually changes in orientation to align across the defect, leading to increased wound strength. In addition collagen type III, which is initially laid down in the immature scar is replaced with the more mature collagen type I.

WOUND DEHISCENCE

Wound dehiscence is disruption of any or all of the layers in a wound. Dehiscence may occur in up to 3 per cent of abdominal wounds. Wound dehiscence most commonly occurs from the 5th to the 8th postoperative day when the strength of the wound is at its weakest. It may herald an underlying abscess and usually presents with a serosanguinous discharge. The patient may have felt a popping sensation during straining or coughing. It is a mechanical wound failure due to various factors causing separation of the closed abdominal wound often with evisceration of the contents.

RISK FACTORS IN WOUND DEHISCENCE

General

- Malnourishment
- Diabetes
- Obesity
- Renal failure
- Jaundice

- Sepsis
- Cancer
- Treatment with steroids

Local

- Inadequate or poor closure of wound
- Poor local wound healing, e.g. because of infection, haematoma or seroma
- Increased intra-abdominal pressure, e.g. in postoperative patients suffering from chronic obstructive airway disease, during excessive coughing

Vertical incision is more prone for dehiscence than transverse as vertical incision cuts across the aponeurosis and when abdominal wall contracts, it creates laterally directed tension on vertical closure. Mass closure is better than layer by-layer closure. Suturing of the peritoneum is not vital in wound failure; it is the fascial closure which decides the strength.

Nonabsorbable monofilament suture material is used for the closure even though evidence says that there is no difference between synthetic absorbable like polyglactic acid and nonabsorbable monofilament suture. Nonabsorbable suture causes prolonged wound pain but is preferred in risk category patients. Suture bite interval should be 1 cm but not more; suture length and wound length ratio should be 4 : 1 or more but not less.

Pinkish serosanguineous discharge (salmon-coloured large quantity of fluid) from the wound. Often omentum or coils of intestine are forced out of the wound. Probing of the wound using gloved finger appreciates dehiscence of musculoaponeurotic layer.

TREATMENT

- Nasogastric aspiration, IV fluids.
- Emergency surgery, i.e. under anaesthesia, wound is opened up properly.
- Coils of intestines are replaced into the abdominal cavity.
- Thorough wash is given. Wound is closed by all layer sutures, passing a nonabsorbable suture material through the red rubber or plastic collar—tension sutures (which is kept for 14 days).
- Antibiotics and IV fluids are continued.
- Wound usually heals well without much second dehiscence. Late problem, may be development of incisional hernia.
- Biological dressings, wound vacuum systems are newer modalities used.

NEWER METHODS OF MANAGING THE BURST ABDOMEN

- Primary deep fascial closure—deep tension/retention sutures
- Delayed closure with initial mesh wrap or pack/ Bogota bag
- Topical negative (vacuum) pressure closure
- Delayed skin closure or skin graft over the dehiscence wound once it granulates well.

- Placing drain in subcutaneous plane
- Suitable antibiotic therapy; fluid management
- Prevention of abdominal compartment syndrome, haematoma or intra-abdominal abscess formation or sepsis.

ANATOMY OF ANTERIOR ABDOMINAL WALL :

The anterior abdominal wall consists of mainly 3 muscles , the External oblique, the Internal oblique and the Transversus abdominis muscles. The 3 muscles form the bulk of the Antero lateral abdominal wall. They give rise anteriorly to broad aponeurosis enclosing the Rectus abdominis muscle forming the Rectus Sheath In the anterior abdominal wall, the external oblique, internal oblique and transversus abdominis muscle join together(condensation of its fascia) to form the linea semilunaris at the outer border of the rectus muscles.

At the midline, the fascia condense again to form the Linea alba . At the level of umbilicus or just below it, the posterior rectus sheath becomes deficient forming the arcuate line or Linea semicircularis. Only the Transversalis fascia covers it posteriorly below the arcuate line.

External Oblique Muscle


Originates Lower 7 ribs

Inserted : Anterior half of the Iliac crest

Direction of Fibres: Superolateral to Inferomedial Direction

The aponeurosis given by the External Oblique muscle passes anterior to the Rectus sheath to insert into the Linea alba. The lower portion of this

aponeurosis folds on itself to form a groove on which the cord structures lie and forms the Inguinal/Poupart's ligament extending from the ASIS to the pubic tubercle.



EXTERNAL OBLIQUE MUSCLE

Internal Oblique Muscle

Originates : 1. Iliopsoas fascia below the lateral half of the Inguinal ligament

2. Anterior 2/3 rd of the Iliac Crest and Lumbodorsal fascia

Inserted:

- The Uppermost fibres insert into the Lower 5 ribs and its cartilages.
- The Central fibers form an aponeurosis at the semilunar fascia and help in the formation of the Rectus sheath
- The lowermost fibers follow an inferomedial course as the same direction as that of the spermatic cord to insert on the pubic tubercle and symphysis pubis.

Direction of fibres : Inferolateral to Superomedial




Fig. 6 - INTERNAL OBLIQUE MUSCLE

Transversus Abdominis muscle

Originates: 1.The lower six costal cartilage

2.spines of the lumbar vertebra

3. Iliac crest

4. Iliopsoas fascia beneath the lateral third of the inguinal ligament.


Inserted:

Direction of fibers and its Insertion :

The muscle fibers traverse in a transverse fashion to give rise to a flat aponeurotic sheet and travels posterior to the rectus abdominis .

The posterior surface of the Transversus abdominis muscle is covered by the Transversalis fascia ,that extends in all directions and forms a complete

Fascial envelope surrounding the abdominal cavity. This fascia is apparently named after the region it covers. It brings together the muscular and aponeurotic fibers into a single layer and supports the weaker areas where the fibers are deficient.



TRANSVERSUS ABDOMINIS MUSCLE

MIDDLE PORTION

Rectus Abdominis and Pyramidal Muscles

The rectus muscle is the master muscle of the abdominal wall.—**Dr. Omar**

The rectus abdominis muscle attaches to the 5th, 6th, and 7th costal cartilages and the xiphoid process. Below, it attaches to the pubic crest, to the ligamentous tissue at the symphysis pubis, and the superior ramus of the pubis

Each rectus muscle is traversed by three tendinous at the level of the xiphoid process, at the umbilicus, and halfway between these points. One or two additional fibrous intersections may occur below the level of the navel. Of these irregular, curved, or zigzagging tendinous bands are usually tightly affixed to the anterior lamina of the rectus sheath. They are occasionally attached to the posterior lamina as well

These fibrous bands attach the rectus muscles firmly to the anterior lamina of the rectus sheath and to the superior attachment to the semirigid thoracic wall. Thus, as the supraumbilical portion of the rectus contracts, that portion of the rectus sheath becomes taut, perhaps assisting in respiratory (or other) physiologic mechanisms.

The rectus muscle is enclosed within a stout sheath formed by the bilaminar aponeuroses of the three flat muscles that divide and pass anteriorly and posteriorly around the muscle, the space between the muscle and the sheath permits the muscle to contract freely with essentially little restraint

. From the rib margin to a point midway between the umbilicus and the pubis (linea semicircularis of Douglas), the posterior sheath is made up of the posterior leaf of the internal oblique aponeurosis, the aponeurosis of the transversus abdominis muscle, and the transversalis fascia. Below this level, the posterior wall is formed by transversalis fascia alone, with variable contributions of aponeurotic bands from the transversus abdominis. The deep epigastric arteries and veins course along the posterior surface of the rectus

muscle, so below the linea semicircularis they are separated from the peritoneum by only transversalis fascia.

The two recti are separated by the linea alba, a tendinous line wherein the three flat muscles both fuse with one another and decussate across the midline. This arrangement is of obvious importance in the contractile properties of the abdominal wall. The linea alba is considerably wider above the umbilicus than below it. Thus, a midline incision inferior to the umbilicus will tend to pass through the laminae of the rectus sheath.

When the supine subject begins to raise the head, the rectus abdominis muscle begins to act before the trunk begins to move. Thus, the rectus abdominis muscle fixes the thorax, so that the sternocleidomastoid muscles can be effective in flexing the neck. Although the rectus is very important in flexing the trunk, it plays little or no role in rotating the trunk. The oblique muscles figure significantly in trunk flexion. The internal oblique is also quite active in maintaining the posture of the upright torso, whereas the external oblique and rectus muscles are quiet.

The internal oblique and transversus abdominis muscles extend superiorly only to the costal margin, whereas the rectus muscle passes ventral to the costal margin in its superior insertion. In this region, therefore, the sternum and costal cartilages provide the posterior wall for the rectus sheath. No aponeurotic lamina is present.

In the lower one-fourth of the abdominal wall, the aponeuroses pass anterior to the rectus muscle as the anterior rectus sheath lamina. The posterior lamina here is formed essentially by transversalis fascia alone together with a highly variable quantity of aponeurotic transversus bundles. This allows passage of the inferior epigastric vascular supply into the sheath.

The pyramidal muscle attaches to the pubic crest and symphyseal ligamentous tissues and inserts into the linea alba. When present, its insertion into the linea alba is a landmark for an accurate midline incision.

The superior and inferior epigastric arteries are responsible for the blood supply of the rectus muscle. The superior epigastric vessels are terminal branches of the internal thoracic artery. The larger, inferior epigastric vessels arise from the external iliacs. The arteries snake behind the muscle within its sheath. The superior and inferior epigastric vessels anastomose at approximately the middle one-third of the muscle.


The most characteristic phenomenon of the superior and inferior epigastric arteries occurs when the muscle contracts: the epigastric vessels glide within their fascial coverings. This avoids injury, bleeding, and the formation of hematoma within the rectus sheath.

Two veins, the superior and inferior epigastric venae comitantes, accompany each epigastric artery.

LINEA ALBA


The linea alba is formed from the xiphoid process of the sternum to the umbilicus. Below the umbilicus its formation is vague and mostly indiscernible. Inferiorly, superficial fibers from the linea alba pass in front of the rectus muscle to find attachment to the symphysis pubis. Deep fibers radiate laterally behind the rectus to attach to the posterior surface of the pubic crest, forming the so-called adminiculum lineae albae.

The subcutaneous tissue of the Anterior abdominal wall is further divided into the Superficial fatty layer of Camper and deep membranous layer of Scarpa which later continues down as the Fascia Lata of the thigh. Including the Scarpa's Fascia during the suturing, helps clear the dead space which is a potential threat for Future wound infections.



The layers of the abdominal wall are :

- Skin
- Subcutaneous tissue
- Superficial fascia
- External oblique muscle
- Internal oblique muscle
- Transversus abdominis muscle
- Transversalis fascia
- Pre- peritoneal adipose and areolar tissue
- Peritoneum.



ANATOMY OF ANTERIOR ABDOMINAL WALL

The subcutaneous tissue of the Anterior abdominal wall is further divided into the Superficial fatty layer of Camper and deep membranous layer of Scarpa which later continues down as the Fascia Lata of the thigh.

Including the Scarpa's Fascia during the suturing, helps clear the dead space which is a potential threat for Future wound infections.

The quality of exposure provided by an incision influences the ease and safety with which an operation can be undertaken and the outcome in ways which defy easy quantification. An incision must provide access to the site of abdominal pathology and allow easy extension if greater exposure than originally anticipated is required. Indeed, the adequacy of an incision is determined above all else by the safety with which an operation can be undertaken.

INCISIONS

The impact that the planning, execution, and closure of an incision has on the outcome of an abdominal operation should not be underestimated. The high combined incidence of surgical site infection (SSI), wound dehiscence, and hernia formation suggests a dominant contribution of wound complications to surgical morbidity. The incisions should provide adequate access to the site of abdominal pathology and should allow extension if greater exposure than originally anticipated is required. Nothing should compromise this and a larger incision or even, on occasion, a second incision should be created without hesitation if exposure is inadequate. Notwithstanding this, the incision should be executed in a fashion that anticipates a secure wound closure and interferes as little as possible with the function and cosmesis of the abdominal wall. These principles apply to both open and laparoscopic incisions. The vertical midline incision remains most popular, and is, perhaps the most flexible.

CHOICE OF INCISION

The relative merit of vertical versus transverse incisions remains a topic of active debate. Proponents of transverse incisions argue that they anticipate a more secure closure than do vertical incisions, a hypothesis supported by anatomic and surgical principle. The fascial fibers of the anterior abdominal wall are oriented transversely or obliquely. Therefore, transverse incisions parallel the direction of the fascial fibers and allow for ready re-approximation with sutures placed perpendicular to these fibers. In contrast, vertical incisions disrupt fascial fibers and must be re-approximated with sutures placed between

fibers. In the latter case, the absence of an anatomic barrier may predispose such sutures to pull through tissue resulting in dehiscence or hernia formation. Despite these concerns, little evidence supports a substantial benefit of transverse incisions.

CLOSURE OF ABDOMINAL INCISIONS

Wound infection is the most common early complication and incisional hernia is the most common long-term complication of open abdominal surgery. Multiple factors contribute to the incidence of wound failure, including diabetes mellitus, malnutrition, obesity, and corticosteroid use. Surgical technique also appears to increase rates of wound failure; however, there has been little consensus regarding the optimal approach to closure.

CLOSURE OF THE FASCIA

The abdomen is closed in multiple layers technique or en mass. The multiple layered technique reconstructs the posterior and anterior aponeurotic sheaths separately with the posterior layer incorporating the peritoneum. Mass closure technique involves a single-layer closure of all the layers and may or may not include the peritoneum.

Given the shorter time required to close the fascial layers en mass, this method is generally preferred. The relative advantages of resorbable versus nonresorbable suture for use in closing the fascia have long been debated. Opponents of closure with nonresorbable suture invoke higher rates of suture sinus formation and increased postoperative pain. In contrast, it has been suggested that closure with resorbable suture may lead to increased incidences

of dehiscence and hernia formation because of intrinsic loss of tensile strength during the postoperative period.

There may be some advantage to the use of slowly resorbable compared to rapidly resorbable suture; the rate of hernia formation when slowly resorbable suture were used is significantly low compared to more rapidly resorbable sutures.

Nonresorbable suture is associated with a higher incidence of suture sinus formation. This association may be greatest with multifilament permanent suture, which may abet bacterial in-growth and infection.

SKIN CLOSURE

A number of skin closure techniques can be used following clean (class I) or clean-contaminated (class II) operations; these include interrupted suture, subcuticular suture, stapled, and adhesive glue. Subcuticular suturing and staples are associated with equivalent rates of wound infection. Glues are used with increasing frequency for skin closure. Advantages of glues include ease and rapidity of application and simplification of wound care; generally, no additional dressing is required. Closure with glues has been compared to traditional skin closure methods in several clinical trials. Wound durability appears to be comparable wound left open if contaminated (class III & IV) and sutured delayed by primary.

RETENTION SUTURES

Incidence of fascial wound dehiscence after major abdominal surgeries is 2–4% and was associated with a mortality rate of 16–22%. Several patient-

related factors are associated with an increased risk of fascial dehiscence, including advanced age, male gender, malnutrition, anemia, and steroids use. However, local mechanical factors and closure technique appear to have a greater influence on the rate of dehiscence. Placement of drains or ostomies through the main incision compromises fascial integrity and should be avoided. Wound sepsis and increased intra-abdominal pressure, whether from ileus, bowel obstruction, atelectasis, or after hernia repair, also compromise the integrity of a fascial closure. Indications for prophylactic placement of retention sutures at initial operation remain controversial. The purpose of retention sutures in this setting is to relieve tension along the suture line in order to prevent significant wound disruption and evisceration in the patient at high risk.

The potential disadvantages of retention sutures, however, are well known and include entrapment of underlying viscera, increased postoperative pain, poor cosmesis, and leakage of intraperitoneal fluid through the wound.

TECHNIQUE OF RETENTION SUTURING

When employed, retention sutures are placed across the wound prior to formal fascial closure. Interrupted permanent mono filament sutures are passed through skin and fascia approximately 2 cm from the wound margin at intervals of several centimeters. Placement is facilitated by the use of a long cutting needle. It may be advantageous to omit the peritoneum from the retention closure in order to protect underlying viscera

from injury or entrapment. After conventional closure of the fascia, the sutures are threaded through rubber tubing bolsters or commercially available plastic bolster devices and tied at the skin level.

TECHNIQUE OF MASS CLOSURE OF THE ABDOMEN

When closing a midline laparotomy incision, two size 0 looped or size 1 nonlooped slowly re-absorbable monofilament sutures are generally used. One suture is anchored at the upper extent and one at the lower extent of the wound. A malleable retractor can be used to protect the underlying viscera while the fascia is closed. The suture is run in a continuous manner, taking full-thickness bites of the linea alba fascia incorporating both the anterior and posterior rectus aponeuroses. Sutures are passed through the fascia a minimum of 1 cm from the wound edge at 1 cm intervals. An assistant holds steady tensions on the suture while the closure progresses. Repetitive relaxation and application of tension of the suture is avoided to limit injury to the fascia. Likewise, it is unnecessary and probably counterproductive to overly tighten the suture as closure progresses, as this may lead to fascial necrosis.



OBSERVATION AND ANALYSIS

This study was done during the period from December 2016 to December 2017 in the department of general surgery, coimbatore Medical college and Hospital, Coimbatore.

The Observations of our study were as follows:

Total number of patients – 60

All patients were operated in emergency surgical theatre, prophylactic retension suturing was done in 30 patients and conventional primary closure was done in 30 patients.


The patients were followed up in post operative period and the status of the wound was observed and the outcome followed was documented in graphs and tables.

RESULTS

The collected data were analysed with IBM.SPSS statics software 21.0 version. To describe about the data descriptive statistics frequency analysis, percentage analysis were used for categorical variables and the mean and S.D were used for continous variables. To find the significant difference between the bivariate samples in Independent groups the Unpaired sample t-test was used. To find the significant categorical data Chi-Square test and Kruskal wallis test was used. In all the above statistical tools the probability value 0.05 is considered as significant


TYPE OF CLOSURE

TYPE OF CLOSURE	NO OF PATIENTS	PERCENTAGE
PRIMARY CLOSURE	30	50%
RETENTION CLOSURE	30	50%




AGE DISTRIBUTION

AGE IN YEARS	NO OF PATIENTS	PERCENTAGE
< 30	12	20%
31-40	10	17%
41-50	12	20%
51-60	15	25%
> 60	11	18%




TYPE OF CLOSURE	AGE IN YEARS	
	MEAN	SD
PRIMARY CLOSURE	38.53	13.65
RETENTION CLOSURE	54.6	12.49
P VALUE - 0.001		
UNPAIRED T TEST		
SIGNIFICANT		




SEX DISTRIBUTION

SEX	NO OF PATIENTS	PERCENTAGE
MALE	46	77%
FEMALE	14	23%




SEX	PRIMARY CLOSURE	RETENTION CLOSURE
MALE	24	22
FEMALE	6	8
P VALUE - 0.542		
CHI SQUARE TEST		
NON SIGNIFICANT		




DIAGNOSIS

DIAGNOSIS	NO OF PATIENTS	PERCENTAGE
APPENDICULAR PERFORATION	3	5%
COLONIC PERFORATION	1	2%
DUODENAL PERFORATION	50	83%
GASTRIC PERFORATION	2	3%
GASTRIC PERFORATION WITH GROWTH	1	2%
ILEAL PERFORATION	3	5%




DIAGNOSIS	PRIMARY CLOSURE	RETENTION CLOSURE
APPENDICULAR PEFORATION	2	1
COLONIC PERFORATION	1	0
DUODENAL PERFORATION	23	27
GASTRIC PERFORATION	1	1
GASTRIC PERFORATION WITH GROWTH	0	1
ILEAL PERFORATION	3	0
P VALUE - 0.341		
KRUSKAL WALLIS TEST		
NON SIGNIFICANT		




PROCEDURE

PROCEDURE	NO OF PATIENTS	PERCENTAGE
APPENDICECTOMY	3	5%
ILEOSTOMY	1	2%
OMENTAL PATCH CLOSURE	50	83%
PRIMARY CLOSURE	3	5%
PRIMARY CLOSURE WITH FJ	1	2%
RESECTION WITH ANASTAMOSIS	2	3%




PROCEDURE	PRIMARY CLOSURE	RETENTION CLOSURE
APPENDICECTOMY	2	1
ILEOSTOMY	1	0
OMENTAL PATCH CLOSURE	23	28
PRIMARY CLOSURE	2	0
PRIMARY CLOSURE WITH FJ	0	1
RESECTION WITH ANASTAMOSIS	2	0
P VALUE - 0.234		
KRUSKAL WALLIS TEST		
NON SIGNIFICANT		




PRE OPERATIVE PERITONITIS DURATION

PRE OPERATIVE PERITONITIS	NO OF PATIENTS	PERCENTAGE
> 24 HRS	53	88%
< 24 HRS	7	12%




PRE OPERATIVE PERITONITIS	PRIMARY CLOSURE	RETENTION CLOSURE
> 24 HRS	26	27
< 24 HRS	4	3
P VALUE - 0.688		
CHI SQUARE TEST		
NON SIGNIFICANT		




EXUDATE


EXUDATE	NO OF PATIENTS	PERCENTAGE
CLEAR	49	82%
PURULENT	9	15%
FECULENT	2	3%




EXUDATE	PRIMARY CLOSURE	RETENTION CLOSURE
CLEAR	19	25
PURULENT	9	3
FECULANT	2	2
P VALUE - 0.001		
KRUSKAL WALLIS TEST		
SIGNIFICANT		




TYPE OF CLOSURE	BLOOD UREA	
	MEAN	SD
PRIMARY CLOSURE	47.1	14.43
RETENTION CLOSURE	39.96	33.37
P VALUE - 0.329		
UNPAIRED T TEST		
NON SIGNIFICANT		




TYPE OF CLOSURE	SERUM CREATININE	
	MEAN	SD
PRIMARY CLOSURE	1.59	0.34
RETENTION CLOSURE	1.47	1.21
P VALUE - 0.679		
UNPAIRED T TEST		
NON SIGNIFICANT		




TYPE OF CLOSURE	HEMOGLOBIN	
	MEAN	SD
PRIMARY CLOSURE	10.68	2.21
RETENTION CLOSURE	9.66	1.62
P VALUE - 0.047		
UNPAIRED T TEST		
SIGNIFICANT		



TYPE OF CLOSURE	TOTAL BILIRUBIN	
	MEAN	SD
PRIMARY CLOSURE	1.89	1.03
RETENTION CLOSURE	2.61	1.38
P VALUE - 0.027		
UNPAIRED T TEST		
SIGNIFICANT		




TYPE OF CLOSURE	SERUM ALBUMIN	
	MEAN	SD
PRIMARY CLOSURE	3.89	1.01
RETENTION CLOSURE	4.08	1.34
P VALUE - 0.536		
UNPAIRED T TEST		
NON SIGNIFICANT		




MANHEIM INDEX

MANHEIM INDEX	NO OF PATIENTS	PERCENTAGE
< 15	39	65%
15-25	19	32%
> 25	2	3%




TYPE OF CLOSURE	MANHEIM INDEX	
	MEAN	SD
PRIMARY CLOSURE	11.66	8.22
RETENTION CLOSURE	9.76	5.47
P VALUE - 0.296		
UNPAIRED T TEST		
NON SIGNIFICANT		




PAIN SCORE

TYPE OF CLOSURE	PAIN SCORE	
	MEAN	SD
PRIMARY CLOSURE	7.23	0.85
RETENTION CLOSURE	4.53	1.07
P VALUE - 0.001		
UNPAIRED T TEST		
SIGNIFICANT		




SEROMA FORMATION

SEROMA FORMATION	NO OF PATIENTS	PERCENTAGE
YES	27	45%
NO	33	55%




SEROMA FORMATION	PRIMARY CLOSURE	RETENTION CLOSURE
PRESENT	18	9
ABSENT	12	21
P VALUE - 0.020		
CHI SQUARE TEST		
NON SIGNIFICANT		




ORGAN FAILURE

ORGAN FAILURE	NO OF PATIENTS	PERCENTAGE
PRESENT	22	37%
ABSENT	38	63%




ORGAN FAILURE	PRIMARY CLOSURE	RETENTION CLOSURE
PRESENT	15	7
ABSENT	15	23
P VALUE - 0.032		
CHI SQUARE TEST		
SIGNIFICANT		




HYPOTENSION

HYPOTENSION	NO OF PATIENTS	PERCENTAGE
PRESENT	21	35%
ABSENT	39	65%




HYPOTENSION	PRIMARY CLOSURE	RETENTION CLOSURE
PRESENT	15	6
ABSENT	15	24
P VALUE - 0.014		
CHI SQUARE TEST		
SIGNIFICANT		




HOSPITAL STAY DAYS

HOSPITAL STAY DAYS	NO OF PATIENTS	PERCENTAGE
< 15 DAYS	23	38%
> 15 DAYS	37	62%




TYPE OF CLOSURE	HOSPITAL STAY IN DAYS	
	MEAN	SD
PRIMARY CLOSURE	22.23	2.4
RETENTION CLOSURE	14.3	2.42
P VALUE - 0.001		
UNPAIRED T TEST		
SIGNIFICANT		




WOUND INFECTION

WOUND INFECTION	NO OF PATIENTS	PERCENTAGE
PRESENT	32	53%
ABSENT	28	47%




WOUND INFECTION	PRIMARY CLOSURE	RETENTION CLOSURE
PRESENT	18	12
ABSENT	14	16
P VALUE - 0.301		
CHI SQUARE TEST		
NON SIGNIFICANT		




WOUND DEHISCENCE

WOUND DEHISCENCE	NO OF PATIENTS	PERCENTAGE
> 24 HRS	17	28%
< 24 HRS	43	72%




WOUND DEHISCENCE	PRIMARY CLOSURE	RETENTION CLOSURE
PRESENT	14	3
ABSENT	16	27
P VALUE - 0.002		
CHI SQUARE TEST		
SIGNIFICANT		




EVISCERATION

EVISCERATION	NO OF PATIENTS	PERCENTAGE
YES	8	13%
NO	52	87%




EVISCERATION	PRIMARY CLOSURE	RETENTION CLOSURE
YES	7	1
NO	23	29
P VALUE - 0.023		
CHI SQUARE TEST		
SIGNIFICANT		




RE-SURGERY

RESURGERY	NO OF PATIENTS	PERCENTAGE
YES	17	28%
NO	43	72%



RESURGERY	PRIMARY CLOSURE	RETENTION CLOSURE
REQUIRED	14	3
NOT REQUIRED	16	27
P VALUE - 0.002		
CHI SQUARE TEST		
SIGNIFICANT		



DISCUSSION

The aim of any surgical procedure is to give relief of symptoms to the patient with minimal or no post operative complications. The recent trend which has been introduced in the field of surgery is to produce the most acceptable surgical scar and numerous studies have been undertaken by implementation of various surgical techniques for the benefit of the patient.

This prospective comparative study was conducted among 60 high risk patients who underwent emergency midline laparotomy for hollow viscus perforation in the emergency general surgery department CMCH, Coimbatore. The study was carried out to compare the efficacy of retention suturing against conventional primary wound closure in midline laparotomies for perforation peritonitis in terms of wound infection, post operative pain, wound dehiscence, hospital stay, re-surgeries.

In our study there were total of 60 (100%) patients, 30 (50%) underwent primary closure and 30 (50%) underwent prophylactic retention suturing for midline wound closure. In this 46 (77%) were males and 14 (23%) were females. The mean age (in years) who underwent primary closure is 38.53 and 54.6 in case of retention closure which is significant ($p = 0.001$).

Post operative pain in the study group was low which was statistically significant ($p = 0.001$). Incidence of wound dehiscence was also low in the study group which was statistically significant ($p = 0.002$). Length of hospital stay (in days) in study group was also low ($p = 0.001$).

One patient in the study group and seven patients in the control group developed evisceration of abdominal contents ($p = 0.023$) which is significant and 3 patients (10.0%) in study group and 14 patients (46.7%) in control group underwent re-surgery (28.3%) ($p = 0.002$).

No statistically significant difference was observed between study and control group in terms of seroma formation and wound infection.

Post-operative morbidity and mortality was found to be significantly low in the study group compared to the control.

Hence prophylactic retention suturing for patients undergoing emergency midline laparotomies for perforation peritonitis is considered as a better choice when compared to primary closure in high risk patients. In spite of all the recent advances in operative technique and risk control method that has been made for wound dehiscence remain high. Surgeons expertise, type of incision made, types of suture material, malnutrition, surgical site infection, prolonged and persistent cough, distension of abdomen, leakage of the pancreatic enzymes, anemia, diabetes, obesity, jaundice, emergency operation, late wound healing due to malignancy all has been a predisposing factors of patients associated with wound dehiscence.

Use of retention suture are one of the recommended technique to reduce the disruption of the fascia in vulnerable cases.

CONCLUSION

Study concludes that Prophylactic Retention suturing in patients with perforation peritonitis undergoing emergency midline laparotomy decreases the incidence of wound dehiscence, reduces pain and lessens hospital stay in high risk patients, when compared with conventional primary wound closure.

BIBLIOGRAPHY

1. Kurata, J.H. Haile, B.M. : Epidemiology of peptic ulcer disease. Clin. Gastroenterol, 13:296, 1984.
2. Kurata, J.H. Henda, G.D., Frank, H. : Hospitalization and mortality rates for peptic ulcers : A comparison of a large health maintenance organization and United States data. Gastroenterology 83: 1008, 2002.
3. Madrazo, B.L., Hricak, II. et al. : Sonographic findings in complicated peptic ulcer, Radiology, 140:457, 1997
4. McGee, G.S., Sawyers J.L. perforated gastric ulcers. Arch. Surg. 122:555, 1987.
5. Miller, R.E., Becker, G.J. Slabsugh, R.D. Detection of pneumoperitoneum: Optimum body position and respiratory phase. AJR.135:487, 1990.
6. Moore, S.W., Fuller, F.W.: Multiple simultaneous complications of peptic ulcer. Am. J. Surg. 97:184, 1989.
7. Rosoff, L., Berne, C.J. : Acute perforation of peptic ulcer, in Nyhus LM, Wastell C. (Eds.): Surgery of the Stomach and Duodenum, 4 edt. Bosten: Little, Brown, 1986, p.466.
8. Sawerys, J.L., Herrington JL, et al. : Acute perforated duodenal ulcer : An evaluation of surgical of surgical management. Arch. Surg. 110 : 527, 1975.

9. Stephen, M. Loewenthal J. : Continuing peritoneal lavage in high risk peritonitis. *Surgery* 85-6036, 1999.
10. Taylor, H. : The non-surgical treatment of perforated peptic ulcer, *Gastroenterology*, 33:353, 1487.
11. Zollinger, R.M, Ayers, H.P. : The Zollinger-Ellison syndrome in Nyhus, L.M. Wastell, C. (eds.) *Surgery of the stomach and Duodenum*, 4 edn Bosten : Little, Brown, 1986, p. 494.
27. Kachroo, R. et al. : Peritonitis (an analysis of 90 cases) *Indian Journal of Surgery*, Vol. 46. April. 1984.
28. Rodney Maingot, *Abdominal*, VOL.1. 9th Edition.
29. Bailey & Love's *Short practice of surgery* .
30. Swaroop M, Williams M, Greene WR et al. Multiple laparotomies are a predictor of fascial dehiscence in the setting of severe trauma. *Am Surg* 2005;71:402–405
31. Gislason H, Grønbech JE, Søreide O. Burst abdomen and incisional hernia after major gastrointestinal operations— comparison of three closure techniques. *Eur J Surg* 1995;161:349–354
32. Penninckx FM, Poelmans SV, Kerremans RP et al. Abdominal wound dehiscence in gastroenterological surgery. *Ann Surg* 1979;189:345–352

33. Pavlidis TE, Galatianos IN, Papaziogas BT et al. Complete dehiscence of the abdominal wound and incriminating factors. *Eur J Surg* 2001;167:351–354
34. Mäkelä JT, Kiviniemi H, Juvonen T et al. Factors influencing wound dehiscence after midline laparotomy. *Am J Surg* 1995;170:387–390
35. Keill RH, Keitzer WF, Nichols WK et al. Abdominal wound dehiscence. *Arch Surg* 1973;106:573–577
36. Col C, Soran A, Col M. Can postoperative abdominal wound dehiscence be predicted? *Tokai J Exp Clin Med* 1998;23:123–127
37. Chin G, Diegelman R, Schultz G: Cellular and molecular regulation of wound healing. In *Wound healing* Edited by: Falabella A, Kirschner R. Boca Raton FL; Taylor, Francis Group; 2005:17-37. Hugh TB: Abdominal wound dehiscence, editorial comment. *Aust NZ J Surgery* 1990, 60:153-155
38. Holmes J.D. Classification of wounds and their management. In: Lumley JSP, Caraven JL. *SurgInt* 1999;45:63-5 A. G. Greenberg and R. P. Saik, “Wound Dehiscence— Pathophysiology and Prevention,” *Archives of Surgery*, Vol. 114, No. 2, 1979, pp. 143-146.

39. Gislason H, Gronbech JE, Soreide O. Burst abdomen and incisional hernia after major gastrointestinal operations. Comparison of three closure techniques. *Eur J Surg* 1995;161:349.
40. Penninckx FM, Poelmans SV, Kerremans RP, et al. Abdominal wound dehiscence in gastroenterological surgery. *Ann Surg* 1979;189:345.
41. Wissing J, van Vroonhoven TJ, Schattenkerk ME, et al. Fascia closure after midline laparotomy: Results of a randomized trial. *Br J Surg* 1987;74:738.
42. Riou JP, Cohen JR, Johnson H Jr. Factors influencing wound dehiscence. *Am J Surg* 1992;163:324.
43. Armstrong CP, Dixon JM, Duffy SW, et al. Wound healing in obstructive jaundice. *Br J Surg* 1984;71:267. 10.Col C, Soran A, Col M. Can postoperative abdominal wound dehiscence be predicted? *Tokai J Exp Clin Med* 1998;23:123.
44. Sørensen LT, Hemmingsen U, Kallehave F, et al. Risk factors for tissue and wound complications in gastrointestinal surgery. *Ann Surg* 2005;241:654.
45. Gislason H, Gronbech JE, Soreide O. Burst abdomen and incisional hernia after major gastrointestinal operations. Comparison of three closure techniques. *Eur J Surg* 1995;161:349.
46. Adams G, Richter RM, Levowitz BS. A safe method of closure with retention sutures. *Surg Gynecol Obstet* 1973;136:981.

47. Banerjee SR, Daoud I, Russell JC, et al. Abdominal wound evisceration. *CurrSurg* 1983;40:432. 15. Miles RM, Moore M, Fitzgerald D, et al. The etiology and prevention of abdominal wound disruption: An analysis of 177 cases. *Am Surg* 1964;30:566.
48. Ponce LC, Morgan MW. A safer wire retention suture. *Am Surg* 1970;36:509.
49. Stivala OG. Retention suture plates. *SurgGynecolObstet* 1983;157:77.
50. Bucknall TE, Cox PJ, Ellis H. Burst abdomen and incisional hernia: A prospective study of 1129 major laparotomies. *Br Med J (Clin Res Ed)* 1982;284:931.
51. Poole GV Jr. Mechanical factors in abdominal wound closure: The prevention of fascial dehiscence. *Surgery* 1985;97:631.
52. Riou JP, Cohen JR, Johnson H Jr. Factors influencing wound dehiscence. *Am J Surg* 1992;163:324.
53. Maäkelaä JT, Kiviniemi H, Juvonen T, et al. Factors influencing wound dehiscence after midline laparotomy. *Am J Surg* 1995; 170:387.
54. Webster C, Neumayer L, Smout R, et al. Prognostic models of abdominal wound dehiscence after laparotomy. *J Surg Res* 2003;109:130.
55. Khan M, Naqvi AH, Irshad K, et al. Frequency and risk factor of abdominal wound dehiscence. *J Coll Physicians Surg Pak* 2004;14:355.
56. Afzal S, Bashir MM. Determinants of wound dehiscence in abdominal surgery in public sector hospital. *Annals of King Edward Medical*

University 2008;14:110. 25. Waldhausen JH, Davies L. Pediatric postoperative abdominal wound dehiscence: Transverse versus vertical incisions. J Am Coll Surg 2000;190:688.

57. Van Ramshorst GH, Nieuwenhuizen J, Hop WCJ, et al. Abdominal wound dehiscence in adults: Development and validation of a risk model. World J Surg 2010;34:20.
58. Rink AD, Goldschmidt D, Dietrich J, et al. Negative side-effects of retention sutures for abdominal wound closure. A prospective randomised study. Eur J Surg 2000; 166:932.
59. Wahl W, Menke H, Schnutgen M, et al. [Fascia dehiscence Cause and prognosis]. Chirurg 1992;63:666.
60. Sørensen LT, Hemmingsen U, Kallehave F, et al. Risk factors for tissue and wound complications in gastrointestinal surgery. Ann Surg 2005;241:654.
61. Reitamo J, Moller C. Abdominal wound dehiscence. Acta Chir Scand 1972;138:170.
62. Richards PC, Balch CM, Aldrete JS. Abdominal wound closure. A randomized prospective study of 571 patients comparing continuous vs. interrupted suture techniques. Ann Surg 1983;197:238.
63. Doughty DB. Preventing and managing surgical wound dehiscence. Adv Skin Wound Care 2005;18:319.

64. Waldrop J, Doughty D. Wound healing physiology. In: Bryant R, editor. Acute and chronic wounds: Nursing management. 2nd ed. St. Louis, MO: Mosby; 2000. p. 17.
65. Posthauer M, Thomas D. Wound care essentials. In: Baranoski S, Ayello E, editors. Nutrition and wound care. Philadelphia: Lippincott: Williams and Wilkins; 2004. p. 379. 35. Carlson MA. Acute wound failure. SurgClin North Am 1997; 77:607.
66. Chavez-Cartaya R, Jiron-Vargas A, Pinto S, et al. Adjustable nylon ties for abdominal wall closure. Am J Surg 1992; 163:609.
67. Boissel P, Jamart J, Grumillier P, et al. A new technique for closing abdominal incisions in patients with poor wound healing. Am J Surg 1982;143:380.
68. Matsuoka J, Gohchi A, Kamikawa Y, et al. Chopstick retention suture for the closure of abdominal wounds. J Am CollSurg 1995;181:471..
69. Heller L, Levin S, Butler C: Management of abdominal wound dehiscence using vacuum assisted closure in patients with compromised healing. Am J Surg 2006, 191:165-172
70. ZhamakKhorgamietal: Prophylactic retention sutures in midline laparotomy in high-risk patients for wound dehiscence.journal homepage: www.JournalofSurgicalResearch.com.
71. Hubbard TB Jr, Rever WB Jr. Retention sutures in the closure of abdominal incisions. Am J Surg 1972;124:378.

72. Goligher JC, Irvin TT, Johnston D, et al. A controlled clinical trial of three methods of closure of laparotomy wounds. *Br J Surg* 1975;62:823.
73. fsraelsson LA, Jonsson T. Overweight and healing of midline incisions. *Eur J Surg* 1997 Mar;163(3) :175-80.
74. Penninckx FM, Poelmans SV, Kerremans RP et a! (1979) Abdominal wound dehiscence in gastroenterological surgery. *Ann Surg* 189:345—352.
75. Zhamak khorgami, Saeed Shoar et al. prophylactic retention sutures in midline laparotomy in high risk cases to prevent wound dehiscence. *Journal of surgical research* 2012:E1-E6
76. Mohanad Abdul Retha. Effect Of Retention Sutures For Prevention Of Abdominal Wound Dehiscence After Laparotomy In High Risk Patients. *iOSR Journal of Pharmacy*;2319-4239.

PROFORMA FOR STUDY ON PERFORATIVE PERITONITIS

DR.HARIDOSS.R

Surgical Unit: II

Coimbatore Medical College

Coimbatore.

NAME:	AGE:	UNIT:	DOA:
ADDRESS:	SEX:	WARD:	DOS:
			DOD:
OCCUPATION:	SOCIOECONOMIC STATUS:		

SOCIO ECONOMICAL STATUS

COMPLAINTS/HISTORY

PAIN

<u>SITE</u>	<u>RADIATION TO</u>	<u>AGRAVATED BY</u>
	<u>RELIEVED BY</u>	
SHOULDER	COUGH	VOMITING
GROIN	MOVEMENT	LYING STILL
LUMBER	FOOD	FOOD
R.LOIN		ANTACIDS
L.LOIN		MILK
NIL		
OTHER		

ONSET PRESENT

PROGRESS
SEVERITY

DURATION

TYPE

NO CHANGE

ONSET

CONTINUOUS

MODERATE

BETTER

PRESENTATION COLICKY

SEVERE

WORSE

SHARP

SYMPTOMS

NAUSEA/VOMITING:

WEIGHT LOSS:

ANOREXIA/DYSPEPSIA

MICTURATION:

FEVER:

BOWELS:

JAUNDICE:

OTHERS:

PAST HISTORY

SIMILAR COMPLAINTS:

DIABETES:

SURGERY:

T.B:

MAJOR ILLNESS:

HYPERTENSION:

OTHERS:

PERSONAL HISTORY

VEG/NON-VEG/MIXED FOOD:

MENSTURATION:

SMOKER:

MARRIED/UNMARRIED:

ALCOHOL:

DRUG INTAKE:

OTHERS:

GENERAL EXAMINATION

BUILT:	TEMPERATURE:
NUTRITION:	PULSE:
HYDRATION:	RESPIRATION:
ANAEMIA:	B.P:
JAUNDICE:	OTHERS:
PEDAL EDEMIA:	

EXAMINATION OF OTHER SYSTEMS

C.V.S:

R.S:

C.N.S:

OTHERS:

EXAMINATION OF ABDOMEN

INSPECTION

DISTENSION:

SCAR:

MOVEMENT WITH RESP:

INJURY:

UMBILICUS:

OTHERS:

PALPATION

TEMP:

MASS:

TENDERNESS:

LIVER:

GUARDING:

SPLEEN:

RIGIDITY:

HERNIA:

OTHERS:

SCRUTUM:

PERCUSSION

LIVER DULLNESS:

OTHERS:

SHIFTING DULLNESS:

FLUID THRILL:

AUSCULTATION: BOWEL SOUNDS:

PERRECTAL EXAMINATION

INVESTIGATION

URINE:

ALBUMIN:

SUGAR:

DEPOSITS:

BLOOD:

H.B %:

T.C.:

D.C.:

SUGAR:

UREA:

WIDAL:

SERUM:

CREATININE:

AMYLASE:

PERITONEAL FLUID CULTURE:

ANALYSIS:

C/S :

RADIOLOGY

PLAIN XRAY – ABDOMEN ERECT:

CHEST P.A.VIEW

ULTRA SOUND ABDOMEN

CT ABDOMEN AND PELVIS

DIAGNOSIS:

TREATMENT

RESUSCITATION

ANALGESIC:

IV F

NASOGASTRIC TUBE

BLADDER CATHETERISATION

ANTIBIOTIC

OTHERS

CONSERVATIVE:

LAPAROTOMY:

DEFINITIVE PROCEDURE:

OTHERS:

FINDINGS:

PERITONEAL FLUID:

SITE OF PERFORATION:

SIZE OF PERFORATION:

ASSOCIATED FINDINGS:

OTHERS:

PROCEDURES:

SIMPLE CLOSURE:

OMENTAL PATCH:

RESECTION/ANASTAMOSIS:

BY PASS:

COLOSTOMY:

SUTURE MATERIALS USED

DRAIN:

POSTOP

NASOGASTRIC SUCTION:

DRAIN REMOVED:

IV F:

SUTURES REMOVED:

INPUT/OUTPUT CHART:

COMPLICATIONS

ANTIBIOTIC:

ANALGESIS/SEDATION:

ON DISCHARGE:

FOLLOW UP:

PROFOMA

Name:

age/sex:

Ip.no.:

Diagnosis:

Procedure done:

Prophylactic retension suturing : done / not done

Parameters:

FIRST LOOK

SECOND LOOK

THIRD LOOK

OUTCOMES:

1.

2.

3.

DISCHARGE DATE:

FOLLOW UP:

CONSENT FORM

I _____, do hereby volunteer and consent to the participate in this study being conducted by Dr. Haridoss. R on “**COMPARATIVE STUDY OF PROPHYLACTIC RETENTION SUTURING VERSUS PRIMARY CLOSURE IN LAPAROTOMIES FOR PERFORATION PERITONITIS AT COIMBATORE MEDICAL COLLEGE HOSPITAL**” . I have read and understood the consent form or it has been read and explained to me in m own native langue in my mother tongue. The stud has been fully explained to me and i may ask questions at any time during the study period.

Signature/Left Thumb impression of the Volunteer

Date:

Place:

Signature and Name of Witness:

Date:

Place:

Signature of the investigator:

Name of the investigator:

ஒப்புதல் படிவம்

நோயாளியின் பெயர் -

தேதி

வயது , பாலினம் -

நேரம்

உள்ளேநோயாளி எண்

நோய்க்குறி

.....

ஆகிய நான் நாளது தேதியில் கோவை மருத்துவக்கல்லூரி
மருத்துவமனையில் அறுவை சிகிச்சைக்காக உள்ளேநோயாளியாக
அனுமதிக்கப்பட்டுள்ளேன்.

எனது இரைப்பை குடலில் உள்ள துளைக்கு அறுவை
செய்யும்போது வயிற்றில் தையல் இடுவதில் வழக்கமான முறை ,
கூடுதல் நிறுத்துதல் தையல் ஆகியவற்றில் உரியதை தெரிவு செய்து
இடப்படுமென மருத்துவர் மூலம் தெரிந்து கொண்டேன்.

எனக்கு உரிய முறையில் அறுவை செய்து கொள்ளவும், எனது
அறுவைக்கு பிந்தைய முடிவுகள் ஆய்வுக்கு உட்படுத்தப்படும்
என்பதையும்,

இந்த ஆய்வு முடிவுகளில் இரகசியம் காக்கப்படும் என்பதையும்,
நான் விரும்பும்போது ஆய்விலிருந்து விலகிக்கொள்ளலாம்
என்பதையும் மருத்துவர் மூலம் தெரிந்து கொண்டேன்.

இவற்றை முழுமையாகப் புரிந்து கொண்டு முழு மனதுடனும்
சுயநினைவுடனும் எவ்வித நிர்பந்தமும் இன்றி சிகிச்சைக்கு ஒத்துழைக்க
ஒப்புக்கொள்கிறேன்.

இடம்

தேதி

நோயாளியின் கையொப்பம்

MASTER CHART

SL.NO.	PATIENT NAME	AGE	SEX	IP.NO.	DIAGNOSIS	PROCEDURE	PREOPERATIVE PERITONITIS > 24 hrs	exudate	hypotensi on	urea	creatinine	organ failure	MAANNHEI N INDEX	PAIN SCORE	Hb	TOTAL BILIRUBI N	ALB UMI N	HOSPITAL STAY (DAYS)	WOUND CLOSURE	SEROMA FORMATION	WOUND INFECTION	WOUND DEHISCENCE	IVISCERATION	RESURGERY
1	RANIDEVI	52	f	207509	duodenal perforation	omental patch closure	y	c	y	56	1.4	y	21	7	9.8	1.9	2.4	21	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
2	AMMAN	22	m	209747	ileal perforation	resection and anastomosis	y	p	y	30	3.8	y	17	6	10.2	2.1	3	26	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
3	MANIKAM	55	m	24598	duodenal perforation	omental patch closure	y	c	n	42	1.2	n	5	4	11	5.2	2.7	18	RETENSION CLOSURE	NO	NO	NO	NO	NO
4	CHINNAMANI	60	f	185980	duodenal perforation	omental patch closure	y	c	n	60	1.8	y	21	5	8.8	1.2	4.2	21	RETENSION CLOSURE	NO	NO	NO	NO	NO
5	DEVI	34	f	219759	gastric perforation	gastric perforation	y	c	y	40	1.2	y	16	4	7.3	2.4	2.4	16	RETENSION CLOSURE	YES	NO	NO	NO	NO
6	RAMGAMMAL	78	f	219148	gastric perforation with growth	primary closure with fj	y	c	y	56	2	y	21	6	11	0.7	3.6	19	RETENSION CLOSURE	NO	NO	NO	NO	NO
7	NIZAM	50	m	218969	duodenal perforation	omental patch closure	y	c	n	36	1	n	9	8	9.7	1.7	2.6	23	PRIMARY CLOSURE	NO	NO	NO	NO	NO
8	KATHIRKAMARAN	50	m	216239	duodenal perforation	omental patch closure	y	c	n	46	0.6	n	9	6	10.2	2.2	4.5	24	PRIMARY CLOSURE	YES	NO	NO	NO	NO
9	ALAGESAN	30	m	215856	duodenal perforation	omental patch closure	y	c	n	33	1.3	n	4	7	11	2.8	4.4	25	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
10	PALANISAMY	55	m	214456	duodenal perforation	omental patch closure	y	c	y	29	0.7	y	16	5	8.9	5.4	5.4	14	RETENSION CLOSURE	YES	YES	YES	NO	YES
11	RANGAN	58	m	213406	duodenal perforation	omental patch closure	y	c	n	30	1.2	n	9	4	9.3	0.8	3.8	18	RETENSION CLOSURE	NO	NO	NO	NO	NO
12	SELVARAJ	52	m	212464	duodenal perforation	omental patch closure	y	c	n	23	0.6	n	9	6	15	1.6	2.7	20	PRIMARY CLOSURE	YES	YES	YES	NO	YES
13	RAJENDRAN	59	m	211397	duodenal perforation	omental patch closure	y	c	y	31	1.2	y	16	3	9	0.7	2.8	15	RETENSION CLOSURE	NO	NO	NO	NO	NO
14	SOLAIAMMAL	29	f	211168	ileal perforation	primary closure	y	p	y	45	1.9	y	22	8	7.7	1.1	4.2	18	PRIMARY CLOSURE	YES	NO	NO	NO	NO
15	RAMATHAL	49	f	210266	duodenal perforation	omental patch closure	n	c	n	26	0.7	n	5	5	10	5.1	5.1	14	RETENSION CLOSURE	NO	NO	NO	NO	NO
16	THANGAVEL	37	m	210234	duodenal perforation	omental patch closure	y	p	y	36	2	y	17	8	9.2	3.1	3.4	23	PRIMARY CLOSURE	NO	YES	YES	NO	YES
17	MURUGAN	48	m	209602	duodenal perforation	omental patch closure	y	c	n	19	1.2	n	4	4	7.6	0.9	3.9	11	RETENSION CLOSURE	NO	NO	NO	NO	NO
18	AALAK	28	m	209383	duodenal perforation	omental patch closure	y	c	y	45	1.1	y	4	7	8.2	1.5	4.4	18	PRIMARY CLOSURE	YES	YES	NO	NO	NO
19	SARASWATHY	65	f	209163	duodenal perforation	omental patch closure	y	c	n	38	1.2	n	14	5	9.9	2.7	5.8	15	RETENSION CLOSURE	YES	NO	NO	NO	NO
20	MANI	57	m	208558	duodenal perforation	omental patch closure	y	c	y	42	1.2	y	16	6	10.2	3.3	3	15	RETENSION CLOSURE	NO	NO	NO	NO	NO
21	SHEBA	16	f	207788	duodenal perforation	omental patch closure	n	c	n	16	0.7	n	5	7	13.1	1.8	3.9	23	PRIMARY CLOSURE	NO	NO	NO	NO	NO
22	KALIDHASS	33	m	204486	duodenal perforation	omental patch closure	y	c	n	32	1	n	4	4	12.4	1.9	4.6	16	RETENSION CLOSURE	NO	NO	NO	NO	NO
23	ELANGO	42	m	200111	duodenal perforation	omental patch closure	y	c	n	25	1.1	n	4	3	11.6	2.5	4.2	13	RETENSION CLOSURE	NO	NO	NO	NO	NO
24	SIVASAMY	58	m	202173	duodenal perforation	omental patch closure	y	c	n	40	0.8	n	9	5	8.2	1	3.7	15	RETENSION CLOSURE	YES	YES	NO	NO	NO
25	MURUGESAN	48	m	201498	duodenal perforation	omental patch closure	n	c	n	26	1.2	n	0	8	8	2.9	2.9	23	PRIMARY CLOSURE	YES	YES	YES	NO	YES
26	RAMACHANDRAN	65	m	198558	duodenal perforation	omental patch closure	y	c	n	32	1.2	n	9	6	7.6	1.6	5.5	12	RETENSION CLOSURE	NO	NO	NO	NO	NO
27	KANNISAMY	47	m	197681	gastric perforation	primary closure	y	c	n	28	1.2	n	4	7	11.1	2.2	3.6	20	PRIMARY CLOSURE	YES	YES	YES	NO	YES
28	MANIKAM	55	m	201598	duodenal perforation	omental patch closure	y	c	n	18	1	n	9	5	9.2	3.6	4.9	14	RETENSION CLOSURE	YES	NO	NO	NO	NO
29	PERUMAL	65	m	195300	duodenal perforation	omental patch closure	y	c	y	39	1.2	n	9	8	6.4	0.9	2.6	21	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
30	RAJESH	38	m	195925	duodenal perforation	omental patch closure	y	p	y	30	0.6	y	17	6	13.4	1.8	5.3	26	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
31	PALANISAMY	64	m	194722	colonic perforation	ileostomy	y	f	y	102	5.2	y	28	7	9.8	2.1	5.8	24	PRIMARY CLOSURE	NO	NO	NO	NO	NO
32	CHARLASS	21	m	192736	duodenal perforation	omental patch closure	y	c	n	46	1.1	n	4	8	12.6	3.6	2.8	19	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES
33	GONAT	61	m	193031	duodenal perforation	omental patch closure	y	c	n	32	0.8	n	9	4	8.8	1.1	6.7	13	RETENSION CLOSURE	YES	YES	YES	NO	YES
34	PALANISAMY	60	m	198715	duodenal perforation	omental patch closure	y	c	n	35	1.1	n	9	6	10.4	4.1	5.8	12	RETENSION CLOSURE	NO	NO	NO	NO	NO
35	CHINNAMANI	60	f	185980	duodenal perforation	omental patch closure	y	c	y	51	3	y	21	7	5.6	1.7	4.7	26	PRIMARY CLOSURE	YES	YES	YES	IVISCERATION	YES

SL.NO.	PATIENT NAME	AGE	SEX	IP.NO.	DIAGNOSIS	PROCEDURE	PREOPERATIVE PERITONITIS > 24 hrs	exudate	hypotensi on	urea	creatinine	organ failure	MANNHEI N INDEX	PAIN SCORE	Hb	TOTAL BILIRUBI N	ALB UMI N	HOSPITAL STAY (DAYS)	WOUND CLOSURE	SEROMA FORMATION	WOUND INFECTION	WOUND DEHISCENCE	OVISCERATION	RESURGERY
36	JESTIN RAJ	37	m	189169	duodenal perforation	omental patch closure	y	p	y	60	0.6	y	17	6	12.2	0.7	4.2	21	PRIMARY CLOSURE	YES	YES	YES	NO	YES
37	PONNUSAMY	65	m	188793	duodenal perforation	omental patch closure	y	c	n	26	1.2	n	9	4	12	2.8	3.6	15	RETENSION CLOSURE	NO	NO	NO	NO	NO
38	KARTHICK	23	m	188596	duodenal perforation	omental patch closure	y	c	n	25	0.8	n	4	8	13.6	3.4	5.5	22	PRIMARY CLOSURE	YES	YES	NO	NO	NO
39	RAMAJAYAM	48	m	187501	duodenal perforation	omental patch closure	y	c	n	30	1.2	n	4	2	9.2	4.6	4.6	14	RETENSION CLOSURE	YES	NO	NO	NO	NO
40	MARTIN	29	m	187080	duodenal perforation	omental patch closure	y	c	n	16	1.2	n	4	9	11	1	4.2	21	PRIMARY CLOSURE	NO	NO	NO	NO	NO
41	JAYAMMAL	65	f	184064	duodenal perforation	omental patch closure	y	c	n	42	1	n	14	5	7	1.9	5.8	14	RETENSION CLOSURE	NO	YES	NO	NO	NO
42	VELLINGIRI	45	m	182212	duodenal perforation	omental patch closure	n	c	n	18	0.6		0	3	8.7	3.4	6.2	11	RETENSION CLOSURE	NO	NO	NO	NO	NO
43	VENUGOPAL	36	m	179792	duodenal perforation	omental patch closure	y	p	y	61	2.4	y	17	7	13	0.8	3.9	21	PRIMARY CLOSURE	YES	YES	YES	NO	YES
44	PRASHANT	23	m	179223	appendicular perforation	appendectomy	n	c	n	23	0.7	n	0	6	11	2.6	4.9	25	PRIMARY CLOSURE	NO	YES	YES	NO	YES
45	PAPPU	23	m	178852	duodenal perforation	omental patch closure	n	c	n	44	1.2	n	0	8	12.4	1.1	3.6	24	PRIMARY CLOSURE	YES	YES	NO	NO	NO
46	PARTHAPPAN	61	m	178505	duodenal perforation	omental patch closure	y	c	n	32	1	n	9	4	11.9	2.4	4.1	12	RETENSION CLOSURE	NO	NO	NO	NO	NO
47	ASHOK	20	m	177964	appendicular perforation	appendectomy	y	p	n	40	2	y	17	7	13.5	3.5	5.5	18	PRIMARY CLOSURE	NO	NO	NO	NO	NO
48	SHANMUGASUNDARAM	45	m	177623	duodenal perforation	omental patch closure	y	c	n	29	1.2	n	4	5	12.6	1.9	4.8	14	RETENSION CLOSURE	YES	NO	NO	NO	NO
49	VIJAYKUMAR	37	m	176476	duodenal perforation	omental patch closure	y	p	y	39	0.6	y	17	6	12	4.6	3.4	21	PRIMARY CLOSURE	NO	NO	NO	NO	NO
50	KANNAN	43	m	176408	duodenal perforation	omental patch closure	y	c	n	35	1.2	n	4	8	10.6	0.7	3.9	25	PRIMARY CLOSURE	NO	NO	NO	NO	NO
51	PICHIMUTHU	75	m	174198	duodenal perforation	omental patch closure	y	c	n	36	1.2	n	9	6	10.7	3.9	4.8	16	RETENSION CLOSURE	NO	NO	NO	NO	NO
52	GURUMOORTHY	42	m	173756	duodenal perforation	omental patch closure	y	p	y	90	1.1	y	17	8	10.9	1.3	5.3	20	PRIMARY CLOSURE	NO	NO	NO	NO	NO
53	PALANIAMMAL	60	f	172803	duodenal perforation	omental patch closure	y	c	n	35	1	n	14	5	7.5	2.1	2.8	12	RETENSION CLOSURE	NO	NO	NO	NO	NO
54	ANGAAPPAN	54	m	16462	ileal perforation	resection and anastomosis	y	f	y	192	5.1	y	28	7	11.3	0.9	2.1	23	PRIMARY CLOSURE	YES	YES	NO	NO	NO
55	AVANTHIKA	2	f	9769	appendicular perforation	appendectomy	n	c	n	26	0.7	n	5	3	8	2.6	2	14	RETENSION CLOSURE	NO	NO	NO	NO	NO
56	KRISHNAMOORTHY	40	m	11987	duodenal perforation	omental patch closure	y	c	n	23	0.7	n	4	6	12	3.4	3.1	12	RETENSION CLOSURE	NO	NO	NO	NO	NO
57	THIRUMALAISAMY	55	m	111537	duodenal perforation	omental patch closure	y	c	n	40	1.2	n	9	4	10	1.8	7.6	11	RETENSION CLOSURE	NO	NO	NO	NO	NO
58	PRIYA 40/F	40	f	10972	duodenal perforation	omental patch closure	y	c	y	60	1.7	y	16	8	9	0.6	4.3	23	PRIMARY CLOSURE	NO	NO	NO	NO	NO
59	BANGASAMY	67	m	5367	duodenal perforation	omental patch closure	y	c	y	92	2.1	y	16	5	9.2	3.4	3.9	13	RETENSION CLOSURE	YES	YES	YES	NO	YES
60	PRIYA	40	F	10972	duodenal perforation	omental patch closure	y	c	n	40	1.2	n	9	8	9	0.6	4.3	23	PRIMARY CLOSURE	NO	NO	NO	NO	NO